

GEORGIAN MEDICAL NEWS

ISSN 1512-0112

№ 9 (330) Сентябрь 2022

ТБИЛИСИ - NEW YORK



ЕЖЕМЕСЯЧНЫЙ НАУЧНЫЙ ЖУРНАЛ

Медицинские новости Грузии
საქართველოს სამედიცინო სიახლენი

GEORGIAN MEDICAL NEWS

Monthly Georgia-US joint scientific journal published both in electronic and paper formats of the Agency of Medical Information of the Georgian Association of Business Press.
Published since 1994. Distributed in NIS, EU and USA.

GMN: Georgian Medical News is peer-reviewed, published monthly journal committed to promoting the science and art of medicine and the betterment of public health, published by the GMN Editorial Board since 1994. GMN carries original scientific articles on medicine, biology and pharmacy, which are of experimental, theoretical and practical character; publishes original research, reviews, commentaries, editorials, essays, medical news, and correspondence in English and Russian.

GMN is indexed in MEDLINE, SCOPUS, PubMed and VINITI Russian Academy of Sciences. The full text content is available through EBSCO databases.

GMN: Медицинские новости Грузии - ежемесячный рецензируемый научный журнал, издаётся Редакционной коллегией с 1994 года на русском и английском языках в целях поддержки медицинской науки и улучшения здравоохранения. В журнале публикуются оригинальные научные статьи в области медицины, биологии и фармации, статьи обзорного характера, научные сообщения, новости медицины и здравоохранения. Журнал индексируется в MEDLINE, отражён в базе данных SCOPUS, PubMed и ВИНТИ РАН. Полнотекстовые статьи журнала доступны через БД EBSCO.

GMN: Georgian Medical News – საქართველოს სამედიცინო სიახლენი – არის ყოველთვიური სამეცნიერო სამედიცინო რეცენზირებადი ჟურნალი, გამოიცემა 1994 წლიდან, წარმოადგენს სარედაქციო კოლეგიისა და აშშ-ის მეცნიერების, განათლების, ინდუსტრიის, ხელოვნებისა და ბუნებისმეტყველების საერთაშორისო აკადემიის ერთობლივ გამოცემას. GMN-ში რუსულ და ინგლისურ ენებზე ქვეყნდება ექსპერიმენტული, თეორიული და პრაქტიკული ხასიათის ორიგინალური სამეცნიერო სტატიები მედიცინის, ბიოლოგიისა და ფარმაციის სფეროში, მიმოხილვითი ხასიათის სტატიები.

ჟურნალი ინდექსირებულია MEDLINE-ის საერთაშორისო სისტემაში, ასახულია SCOPUS-ის, PubMed-ის და ВИНТИ РАН-ის მონაცემთა ბაზებში. სტატიების სრული ტექსტი ხელმისაწვდომია EBSCO-ს მონაცემთა ბაზებიდან.

WEBSITE

www.geomednews.com

К СВЕДЕНИЮ АВТОРОВ!

При направлении статьи в редакцию необходимо соблюдать следующие правила:

1. Статья должна быть представлена в двух экземплярах, на русском или английском языках, напечатанная через **полтора интервала на одной стороне стандартного листа с шириной левого поля в три сантиметра**. Используемый компьютерный шрифт для текста на русском и английском языках - **Times New Roman (Кириллица)**, для текста на грузинском языке следует использовать **AcadNusx**. Размер шрифта - **12**. К рукописи, напечатанной на компьютере, должен быть приложен CD со статьей.

2. Размер статьи должен быть не менее десяти и не более двадцати страниц машинописи, включая указатель литературы и резюме на английском, русском и грузинском языках.

3. В статье должны быть освещены актуальность данного материала, методы и результаты исследования и их обсуждение.

При представлении в печать научных экспериментальных работ авторы должны указывать вид и количество экспериментальных животных, применявшиеся методы обезболивания и усыпления (в ходе острых опытов).

4. К статье должны быть приложены краткое (на полстраницы) резюме на английском, русском и грузинском языках (включающее следующие разделы: цель исследования, материал и методы, результаты и заключение) и список ключевых слов (key words).

5. Таблицы необходимо представлять в печатной форме. Фотокопии не принимаются. **Все цифровые, итоговые и процентные данные в таблицах должны соответствовать таковым в тексте статьи**. Таблицы и графики должны быть озаглавлены.

6. Фотографии должны быть контрастными, фотокопии с рентгенограмм - в позитивном изображении. Рисунки, чертежи и диаграммы следует озаглавить, пронумеровать и вставить в соответствующее место текста **в tiff формате**.

В подписях к микрофотографиям следует указывать степень увеличения через окуляр или объектив и метод окраски или импрегнации срезов.

7. Фамилии отечественных авторов приводятся в оригинальной транскрипции.

8. При оформлении и направлении статей в журнал МНГ просим авторов соблюдать правила, изложенные в «Единых требованиях к рукописям, представляемым в биомедицинские журналы», принятых Международным комитетом редакторов медицинских журналов - <http://www.spinesurgery.ru/files/publish.pdf> и http://www.nlm.nih.gov/bsd/uniform_requirements.html В конце каждой оригинальной статьи приводится библиографический список. В список литературы включаются все материалы, на которые имеются ссылки в тексте. Список составляется в алфавитном порядке и нумеруется. Литературный источник приводится на языке оригинала. В списке литературы сначала приводятся работы, написанные знаками грузинского алфавита, затем кириллицей и латиницей. Ссылки на цитируемые работы в тексте статьи даются в квадратных скобках в виде номера, соответствующего номеру данной работы в списке литературы. Большинство цитированных источников должны быть за последние 5-7 лет.

9. Для получения права на публикацию статья должна иметь от руководителя работы или учреждения визу и сопроводительное отношение, написанные или напечатанные на бланке и заверенные подписью и печатью.

10. В конце статьи должны быть подписи всех авторов, полностью приведены их фамилии, имена и отчества, указаны служебный и домашний номера телефонов и адреса или иные координаты. Количество авторов (соавторов) не должно превышать пяти человек.

11. Редакция оставляет за собой право сокращать и исправлять статьи. Корректур авторам не высылаются, вся работа и сверка проводится по авторскому оригиналу.

12. Недопустимо направление в редакцию работ, представленных к печати в иных издательствах или опубликованных в других изданиях.

При нарушении указанных правил статьи не рассматриваются.

REQUIREMENTS

Please note, materials submitted to the Editorial Office Staff are supposed to meet the following requirements:

1. Articles must be provided with a double copy, in English or Russian languages and typed or computer-printed on a single side of standard typing paper, with the left margin of 3 centimeters width, and 1.5 spacing between the lines, typeface - **Times New Roman (Cyrillic)**, print size - 12 (referring to Georgian and Russian materials). With computer-printed texts please enclose a CD carrying the same file titled with Latin symbols.

2. Size of the article, including index and resume in English, Russian and Georgian languages must be at least 10 pages and not exceed the limit of 20 pages of typed or computer-printed text.

3. Submitted material must include a coverage of a topical subject, research methods, results, and review.

Authors of the scientific-research works must indicate the number of experimental biological species drawn in, list the employed methods of anesthetization and soporific means used during acute tests.

4. Articles must have a short (half page) abstract in English, Russian and Georgian (including the following sections: aim of study, material and methods, results and conclusions) and a list of key words.

5. Tables must be presented in an original typed or computer-printed form, instead of a photocopied version. **Numbers, totals, percentile data on the tables must coincide with those in the texts of the articles.** Tables and graphs must be headed.

6. Photographs are required to be contrasted and must be submitted with doubles. Please number each photograph with a pencil on its back, indicate author's name, title of the article (short version), and mark out its top and bottom parts. Drawings must be accurate, drafts and diagrams drawn in Indian ink (or black ink). Photocopies of the X-ray photographs must be presented in a positive image in **tiff format**.

Accurately numbered subtitles for each illustration must be listed on a separate sheet of paper. In the subtitles for the microphotographs please indicate the ocular and objective lens magnification power, method of coloring or impregnation of the microscopic sections (preparations).

7. Please indicate last names, first and middle initials of the native authors, present names and initials of the foreign authors in the transcription of the original language, enclose in parenthesis corresponding number under which the author is listed in the reference materials.

8. Please follow guidance offered to authors by The International Committee of Medical Journal Editors guidance in its Uniform Requirements for Manuscripts Submitted to Biomedical Journals publication available online at: http://www.nlm.nih.gov/bsd/uniform_requirements.html
http://www.icmje.org/urm_full.pdf

In GMN style for each work cited in the text, a bibliographic reference is given, and this is located at the end of the article under the title "References". All references cited in the text must be listed. The list of references should be arranged alphabetically and then numbered. References are numbered in the text [numbers in square brackets] and in the reference list and numbers are repeated throughout the text as needed. The bibliographic description is given in the language of publication (citations in Georgian script are followed by Cyrillic and Latin).

9. To obtain the rights of publication articles must be accompanied by a visa from the project instructor or the establishment, where the work has been performed, and a reference letter, both written or typed on a special signed form, certified by a stamp or a seal.

10. Articles must be signed by all of the authors at the end, and they must be provided with a list of full names, office and home phone numbers and addresses or other non-office locations where the authors could be reached. The number of the authors (co-authors) must not exceed the limit of 5 people.

11. Editorial Staff reserves the rights to cut down in size and correct the articles. Proof-sheets are not sent out to the authors. The entire editorial and collation work is performed according to the author's original text.

12. Sending in the works that have already been assigned to the press by other Editorial Staffs or have been printed by other publishers is not permissible.

**Articles that Fail to Meet the Aforementioned
Requirements are not Assigned to be Reviewed.**

ავტორთა საქურაღებოლ!

რედაქციაში სტატიის წარმოდგენისას საჭიროა დაიცვათ შემდეგი წესები:

1. სტატია უნდა წარმოადგინოთ 2 ცალად, რუსულ ან ინგლისურ ენებზე დაბეჭდილი სტანდარტული ფურცლის 1 გვერდზე, 3 სმ სიგანის მარცხენა ველისა და სტრიქონებს შორის 1,5 ინტერვალის დაცვით. გამოყენებული კომპიუტერული შრიფტი რუსულ და ინგლისურენოვან ტექსტებში - **Times New Roman (Кириллица)**, ხოლო ქართულენოვან ტექსტში საჭიროა გამოვიყენოთ **AcadNusx**. შრიფტის ზომა – 12. სტატიას თან უნდა ახლდეს CD სტატიით.

2. სტატიის მოცულობა არ უნდა შეადგენდეს 10 გვერდზე ნაკლებს და 20 გვერდზე მეტს ლიტერატურის სიის და რეზიუმეების (ინგლისურ, რუსულ და ქართულ ენებზე) ჩათვლით.

3. სტატიაში საჭიროა გაშუქდეს: საკითხის აქტუალობა; კვლევის მიზანი; საკვლევი მასალა და გამოყენებული მეთოდები; მიღებული შედეგები და მათი განსჯა. ექსპერიმენტული ხასიათის სტატიების წარმოდგენისას ავტორებმა უნდა მიუთითონ საექსპერიმენტო ცხოველების სახეობა და რაოდენობა; გაუტკივარებისა და დაძინების მეთოდები (მწვავე ცდების პირობებში).

4. სტატიას თან უნდა ახლდეს რეზიუმე ინგლისურ, რუსულ და ქართულ ენებზე არანაკლებ ნახევარი გვერდის მოცულობისა (სათაურის, ავტორების, დაწესებულების მითითებით და უნდა შეიცავდეს შემდეგ განყოფილებებს: მიზანი, მასალა და მეთოდები, შედეგები და დასკვნები; ტექსტუალური ნაწილი არ უნდა იყოს 15 სტრიქონზე ნაკლები) და საკვანძო სიტყვების ჩამონათვალი (key words).

5. ცხრილები საჭიროა წარმოადგინოთ ნაბეჭდი სახით. ყველა ციფრული, შემაჯამებელი და პროცენტული მონაცემები უნდა შეესაბამებოდეს ტექსტში მოყვანილს.

6. ფოტოსურათები უნდა იყოს კონტრასტული; სურათები, ნახაზები, დიაგრამები - დასათაურებული, დანომრილი და სათანადო ადგილას ჩასმული. რენტგენოგრაფიების ფოტოასლები წარმოადგინეთ პოზიტიური გამოსახულებით **tiff** ფორმატში. მიკროფოტოსურათების წარწერებში საჭიროა მიუთითოთ ოკულარის ან ობიექტივის საშუალებით გადიდების ხარისხი, ანათალების შედეგების ან იმპრეგნაციის მეთოდი და აღნიშნოთ სურათის ზედა და ქვედა ნაწილები.

7. სამამულო ავტორების გვარები სტატიაში აღინიშნება ინიციალების თანდართვით, უცხოურისა – უცხოური ტრანსკრიპციით.

8. სტატიას თან უნდა ახლდეს ავტორის მიერ გამოყენებული სამამულო და უცხოური შრომების ბიბლიოგრაფიული სია (ბოლო 5-8 წლის სიღრმით). ანბანური წყობით წარმოდგენილ ბიბლიოგრაფიულ სიაში მიუთითეთ ჯერ სამამულო, შემდეგ უცხოელი ავტორები (გვარი, ინიციალები, სტატიის სათაური, ჟურნალის დასახელება, გამოცემის ადგილი, წელი, ჟურნალის №, პირველი და ბოლო გვერდები). მონოგრაფიის შემთხვევაში მიუთითეთ გამოცემის წელი, ადგილი და გვერდების საერთო რაოდენობა. ტექსტში კვადრატულ ფხიხლებში უნდა მიუთითოთ ავტორის შესაბამისი N ლიტერატურის სიის მიხედვით. მიზანშეწონილია, რომ ციტირებული წყაროების უმეტესი ნაწილი იყოს 5-6 წლის სიღრმის.

9. სტატიას თან უნდა ახლდეს: ა) დაწესებულების ან სამეცნიერო ხელმძღვანელის წარდგინება, დამოწმებული ხელმოწერითა და ბეჭდით; ბ) დარგის სპეციალისტის დამოწმებული რეცენზია, რომელშიც მითითებული იქნება საკითხის აქტუალობა, მასალის საკმაობა, მეთოდის სანდოობა, შედეგების სამეცნიერო-პრაქტიკული მნიშვნელობა.

10. სტატიის ბოლოს საჭიროა ყველა ავტორის ხელმოწერა, რომელთა რაოდენობა არ უნდა აღემატებოდეს 5-ს.

11. რედაქცია იტოვებს უფლებას შეასწოროს სტატია. ტექსტზე მუშაობა და შეჯერება ხდება საავტორო ორიგინალის მიხედვით.

12. დაუშვებელია რედაქციაში ისეთი სტატიის წარდგენა, რომელიც დასაბეჭდად წარდგენილი იყო სხვა რედაქციაში ან გამოქვეყნებული იყო სხვა გამოცემებში.

აღნიშნული წესების დარღვევის შემთხვევაში სტატიები არ განიხილება.

Содержание:

P.V. Fedorych, T.V. Kuts, S.B. Koval. DETERMINATION OF THE SENSITIVITY OF GARDNERELLA VAGINALIS TO FENTICONAZOLE.....	6-10
Giuseppe Taccardo, Andrea Perna, Alessandro Domenico Santagada, Marco Passiatore, Calogero Velluto, et al. DOES AN EARLY POST-OPERATIVE PAIN RELIEVE INFLUENCE THE FUNCTIONAL OUTCOME OF PATIENTS WITH COLLES FRACTURES TREATED WITH EG-BLOCK SYSTEM?.....	11-16
Oksana Knyzhenko, Svitlana Knyzhenko, Krainyk Hryhorii, Kseniia Kotlubaieva. IMPROPER PERFORMANCE OF PROFESSIONAL DUTIES BY A MEDICAL PROFESSIONAL: CURRENT ISSUES OF RESPONSE AND INVESTIGATION UNDER CRIMINAL LAW.....	17-22
Fana Lichoska-Josifovikj, Kalina Grivceva-Stardelova, Beti Todorovska, Vladimir Andreevski, Filip Nikolov, Dzem Adem. THE VALUE OF SERUM-ASCITES ALBUMIN GRADIENT AS A PREDICTOR OF SPONTANEOUS BACTERIAL PERITONITIS IN PATIENTS WITH LIVER CIRRHOSIS AND ASCITES.....	23-25
Mher S. Bisharyan, Ara B. Dallakyan. ASSESSMENT OF THE SOCIAL AND MEDICAL ASPECTS OF SUICIDE IN THE REPUBLIC OF ARMENIA.....	26-31
Nadiya Ye. Barabash, Tetiana M. Tykhonova, Diana M. Dorosh, Larysa O. Martymianova. HETEROGENEITY OF CLINICAL MANIFESTATIONS OF HYPERPROLACTINEMIA (REVIEW AND OWN OBSERVATIONS)	32-36
Alexander Schuh, Philipp Koehl, Stefan Sesselmann, Tarun Goyal, Achim Benditz. INCIDENTAL INTRAOSSEOUS CALCANEAL LIPOMA IN A PATIENT SUFFERING FROM PLANTARFASZIITIS	37-39
Alexander Schuh, Philipp Koehl, Stefan Sesselmann, Tarun Goyal, Achim Benditz. INTRAMUSCULAR MYXOMA OF THE BUTTOCK- A CASE REPORT	40-42
Tsvetkova M. A., Kovalenko A. YU. ALGORITHM OF ORTHODONTIC TREATMENT PATIENTS WITH A BURDENED DRUG ANAMNESIS. DRUGS THAT CAN INHIBIT TOOTHMOVEMENT.....	43-48
Mazin M. Hammady, Shaymaa J. Mohammed. IMPLEMENTING NEW TECHNIQUE TO EVALUATE COGNITIVE FUNCTION IN PATIENTS WITH MIGRAINE DURING THE ATTACK.....	49-53
Nataliia O. Shevchenko, Liliya S. Babinets, Iryna M. Halabitska. AGE-DEPENDENT IMMUNE STATUS CHANGES IN CHRONIC PANCREATITIS PATIENTS.....	54-58
Salah Kadhim Muslim. A SINGLE SURGEON'S EXPERIENCE IN DEFINING THE LEARNING CURVE FOR TRANSORAL ENDOSCOPIC THYROIDECTOMY -VESTIBULAR APPROACH (TOETVA).....	59-64
Muradyan A.E. CORRELATION AND INFRASTRUCTURE OF SOME PHYSICAL HEALTH INDICATORS BEFORE AND DURING COVID-19 PANDEMIC.....	65-69
Brych V.V., Vasylynets M.M., Shmanko O.P., Bilak-Lukyanchuk V.Y PARTICIPATION OF TRAUMATOLOGISTS IN PROVIDING MEDICAL REHABILITATION OF PATIENTS WITH INJURIES AT THE REGIONALLEVEL.....	70-73
Soldatiuk V.M., Rozhko M.M., Pantus A.V CLINICAL-MORPHOLOGICAL SUBSTANTIATION OF THE FIBROUS MATRIX WITH BIOGEL CENO BONETM APPLICATION FOR PRESERVATION OF THE ALVEOLAR PROCESS OF THE JAWS AFTER THE TEETH REMOVAL.....	74-80
O. Rotar, I. Khomiak, R. Sydorchuk, S. Boiko, I. Bilyk, O. Hrama, Y. Migaichuk. EFFICACY OF THE ALGORITHMIC STEP-UP APPROACH OF INTERVENTIONAL TREATMENT OF PATIENTS WITH ACUTE NECROTIZINGPANCREATITIS.....	81-85
V.V. Ohorenko, A.V. Shornikov, A.G. Kyrychenko, Y.N. Zavalko, V.N. Khomyakov, N.V. Tomakh. IMPROVEMENT OF QUALITY OF LIFE FOR PATIENTS WITH ASEPTIC NECROSIS OF THE FEMORAL HEAD AND NON- PSYCHOTIC MENTAL DISORDERS.....	86-89
Nigar Karimova Ildirim. CYP2B6 SINGLE NUCLEOTIDE POLYMORPHISMS IN AN AZERBAIJANI POPULATION.....	90-93
Olha Filyk, Yaroslav Pidhirnyi. RESPIRATORY MUSCLES FUNCTION IN CHILDREN 6-18 YEARS OLD WITH ACUTE HYPOXEMIC RESPIRATORY FAILURE: THE PROSPECTIVE OBSERVATIONAL COHORT STUDY.....	94-98

Héctor M. Ramos-Zaldívar, Karla G. Reyes-Perdomo, Nelson A. Espinoza-Moreno, Ernesto Tomás Dox-Cruza, Thania Camila Aguirre Urbinaa, et al.	
SAFETY AND EFFICACY OF THYMIC PEPTIDES IN THE TREATMENT OF HOSPITALIZED COVID-19 PATIENTS IN HONDURAS.....	99-105
Melnychenko MH, Kvashnina AA, Sytnikova VA.	
PROGNOSTIC MODEL OF POSTOPERATIVE ADHESIVE INTESTINAL OBSTRUCTION RISK IN CHILDREN.....	106-109
Musayev SA.	
EVALUATION OF THE QUALITY OF LIFE AFTER REVASCULARIZATION AND RECONSTRUCTIVE OPERATIONS ON MITRAL VALVE IN PATIENTS WITH CORONARY HEART DISEASE.....	110-114

PROGNOSTIC MODEL OF POSTOPERATIVE ADHESIVE INTESTINAL OBSTRUCTION RISK IN CHILDREN

Melnychenko MH, Kvashnina AA, Sytnikova VA.

The Odessa National Medical University, Odessa, Ukraine.

Abstract.

Objective: to develop a prognostic model of the postoperative adhesive intestinal obstruction risk in children by the use of multivariate statistics.

Materials and methods: To create a model for the prediction of the postoperative adhesive intestinal obstruction by using the method of binary logistic regression, a retrospective analysis of the features of the acute inflammatory disease of the abdominal cavity organs and treatment strategy in 119 children.

Results: The level of significance of clinical and anamnestic signs in children with appendicular peritonitis and the postoperative adhesive intestinal obstruction was determined using Pearson's test χ^2 . The main factors (leukopenia at hospitalization) ($p=0.033$) pronounced stigmatization with signs of connective tissue dysplasia ($p=0.013$) and duration of surgery ($p=0.0002$) were highlighted, which indicate the postoperative adhesive intestinal obstruction risk. The logistic regression method was used to create a prognostic model for calculating the postoperative adhesive intestinal obstruction risk, which was assessed as "very good" by ROC analysis (test sensitivity – 77%; specificity – 72%); AUC = 0.796.

Conclusions: The application of the proposed prognostic model of the postoperative adhesive intestinal obstruction risk in children with acute inflammatory diseases of the abdominal organs allows to make decision as for the extent of surgery and optimization of treatment strategy at all stages of rendering care.

Key words. Prognostic model, risk, postoperative adhesive intestinal obstruction, children.

Introduction.

Determining the risk of intra-abdominal adhesions in children with acute abdominal pathology helps to optimize the treatment of such patients and prevent the development of postoperative adhesive intestinal obstruction (PAIO), which occurs in 62% of cases [1-4]. Since peritoneal adhesiogenesis is multifactorial, the use of individual clinical and anamnestic factors or their set without considering the total effect for a concrete patient does not allow to reliably predict the development of adhesions [2,5,6]. The best way to diagnose the PAIO risk in children, in our opinion, is to develop a prognostic model by using multivariate statistics to identify the mutual influence of predictors to optimize preventive measures.

Materials and methods.

To create a method for predicting PAIO by the way of the binary logistic regression, a retrospective analysis of the clinical course of acute inflammatory disease of the abdominal organs (AO) and treatment strategy was performed in 119 children who were treated at the surgical departments of the Regional Children's Clinical Hospital in Odessa. The first group (A) consisted of 59 children with adhesive intestinal obstruction (AIO) occurred

after surgical treatment of appendicular peritonitis (AP). The second group (B) included 60 children with AP who had no signs of AIO during the postoperative follow-up period.

To determine the logistic regression parameters, the selection of predictors with the known AIO in both groups of children was conducted. The logistic function looks like this:

$$f(z) = \frac{1}{1 + e^{-z}}$$

$$z(x) = \beta_0 + \beta_1 x_1 + \dots + \beta_n x_n$$

where: f – gets the value between 0 и 1; β_0, \dots, β_n – logistic regression parameters. x_1, \dots, x_n – signs determination.

This equation allows us to estimate the probability of the event (in our case PAIO) in each study participant with an individual set of predictors. For significant assessment of the diagnostic test, the area under the characteristic curve (AUC) is determined, which reflects the optimal value of sensitivity and specificity of each feature [7-10]. Statistical processing of the study results was performed using MedCalc 9.03 packages: Statistica 8.0 and SPSS 11.01.

Results.

The first step in creating a diagnostic model was to identify statistically significant factors having influence on the PAIO risk. With this purpose, we analyzed the diagnostic value of 47 signs, which were considered by potential predictors: demographic and individual factors of the patient, clinical and anamnestic signs, laboratory parameters, intraoperative data and extent of surgery, features of the postoperative period course, the presence and severity of undifferentiated connective tissue dysplasia (UCTD). In addition, the acetylation genotype was studied by determining point mutations in the N-acetyltransferase 2 (NAT2) gene using polymerase chain reaction and restriction fragment length analysis. The list of features that were analyzed is shown in table 1.

Relevant features were identified by filtering low-frequency features. In our case, the features that in 90% or more cases take the same value were excluded, because they do not improve the prognostic ability of the model. At that stage, 12 binary features were excluded.

The next step for the remained signs a statistical significance to assess the risk of PAIO by using Pearson's criterion χ^2 and ANOVA variance analysis was calculated. Empirically, the features with the lowest p-value were selected to create a logistic model. Simultaneously the factors were identified that demonstrated a statistically significant relationship with the PAIO development: leukopenia at the time of hospitalization ($p = 0.033$), pronounced stigmatization of NCTD signs ($p = 0.013$), duration of surgery ($p = 0.0002$).

Table 1. Potential predictors of the adhesive intestinal obstruction risk.

Demographic/ individual	age, sex; place of residence (city/countryside); belonging to socially and economically disadvantaged groups; <i>NAT2</i> acetylation genotype; presence and degree of NCTD; functional disorders of the digestive tract in the anamnesis
Clinical and anamnestic signs	The duration of the disease at the time of treatment; general condition at hospitalization; degree of exsiccosis; localization and nature of pain, defense, peritoneal signs; the nature of vomiting; violation of defecation and gas discharge; fever; rectal examination data.
Laboratory indicators	The presence of hyperleukocytosis; the presence of leukopenia; total number of leukocytes, leukocyte shift; hematocrit; C-reactive protein; fibrinogen; dyselectrolytemia; acid-base condition; serum glucose.
Intraoperative data and scope of intervention	Type of operational access; type of peritonitis; position of the appendix; the presence and nature of effusion; the presence of fibrin in the abdominal cavity; technical difficulties; omentectomy; intestinal deserosation; drainage; duration of intervention.
Features of the postoperative period	The presence of complications (intra-abdominal / from the wound); relaparotomy; restoration of peristalsis / independent defecation; removal of nasogastric probe; the beginning of enteral nutrition (a day); prokinetics stimulation; duration of stay at the intensive care unit and inpatient treatment

Table 2. Logistic Regression Equation Predictors.

	Predictors	β Coefficient
x_1	pronounced stigmas of NCTD	1.031
x_2	midline laparotomy	0.710
x_3	leukopenia at hospitalization	0.697
x_4	intra-abdominal complications at the early postoperative period	0.475
x_5	duration of intervention, min	0.027
x_6	local unrestricted peritonitis	-0.516
x_7	no vomiting	-0.755
x_8	medial position of the appendix	-1.004

According to the results of the analysis, surgical treatment of peritonitis is one of the most important risk factors for adhesions. The performance of omentectomy and the total intervention time are the factors that showed the lowest p-value, and accordingly have a significant impact on the postoperative adhesion process. In particular, among children with AIO, omentectomy was performed in 21 (35.6%) cases, in group B in 41 (68.3%) cases. Pearson’s criterion χ^2 is 12.78; $p < 0.001$. These indicators also correlate with our observations as for intraoperative data in the surgical treatment of AIO, because in a significant number of cases the cause of obstruction is joints involving a strand of a large omentum or the omentum itself, soldered to the postoperative scar. In addition, omentitis, intra-abdominal infiltrates and omental abscesses are the main causes of relaparotomy at the early postoperative period of peritonitis treatment in children, which are independent predictors of excessive adhesions risk.

The duration of surgery has a significant impact on the PAIO risk. For children from the main group, this figure averaged (77.7 ± 23.99) minutes, in the control group – (64.0 ± 13.74) minutes. It should be noted that this indicator is integral and depends, for example, both on the experience of the surgeon and the clinical situation, including the location of the inflammatory focus, technical difficulties, the occurrence of intraoperative complications, the spreading of intra-abdominal inflammation etc.

The last step is to analyze all possible combinations of selected features. For each combination, logistic regression was studied and evaluated by cross-checking using AUC ROC analysis, as this indicator reflects the combination of sensitivity and specificity of the proposed predicting model [8-10].

The obtained logistic regression equation includes 8 predictors and corresponding coefficients (Table 2).

The value of the free member of the equation β_0 was (-1.5193). So, a probability of PAIO development, namely the value (z) in the logistic regression equation is calculated as follows:

$$z(x) = -1.5193 + 1.031x_1 + 0.810x_2 + 0.697x_3 + \dots - 1.004x_8$$

As this formula demonstrates, the negative value of the coefficient means the protective effect of the sign on PAIO development. Calculation of the PAIO developing probability $-f(z)$ according to this formula, classifies the child’s belonging to the PAIO risk group. With the purpose of automatical calculation of this indicator using the mathematical or statistical analysis program, we have created an appropriate algorithm in Microsoft Office Excel.

The quality assessment of the prognostic model was performed by cross-checking using AUC ROC-analysis. Figure 1 shows the characteristic curve by values $f(z)$. The optimal dividing point corresponds to the value 0.0.




Figure 1. Characteristic curve of the prognostic model ROC-analysis ($Sp = 77\%$, $Se = 72\%$).




Figure 2. Example of calculating the PAIO risk using MO Excel.

If the patient has $f(z) \geq 0.5$, the model is considered positive for the risk of PAIO developing according to certain predictors. The area under the ROC curve was AUC 0.796, which indicated a “very good” operation of the prognostic model.

The prognostic specificity of this model is (Sp) = 77%, sensitivity (Se) = 72%. The probability of intraperitoneal adhesions is 72% (test sensitivity). The statistical estimation of the area under the ROC curve is quite high ($p = 0.0001$), so developed on the basis of the logistic regression, the proposed model can be used to prognosis of the PAIO risk in children with acute inflammatory diseases and choice of an adequate treatment strategy.

Here is an example of calculating the PAIO risk developing in a child T. of 7 years old with appendicular peritonitis (Medical card No 90548. Postoperative diagnosis: Gangrenous-perforated appendicitis, widespread peritonitis) based on the results of entering the values of the corresponding predictors in Figure 2.

According to the calculations of the logistic regression equation, the child belongs to the PAIO risk group, because $f(z) = 0.5773$, i.e., $f(z)$ is closure to 1. In 11 months after the initial intervention, the child was hospitalized with signs of acute intestinal obstruction. During the operation, an extensive adhesion process in the abdominal cavity, multiple visceroparietal and viscerovisceral adhesions with the formation of “double-barrels” were determined. Obstruction was caused by a cord-like adhesion between the loop of the small intestine and the anterior abdominal wall, which caused strangulation of the adjacent loop of the intestine.

So, the prognostic model of the PAIO risk by logistic regression showed a good result, and the predictor that showed the greatest connection with the development of AIO is the duration of abdominal surgery. The application of the proposed prognostic model of the PAIO risk by logistic regression method of selected signs at the stage of clinical history and intraoperative data allowed to make a decision as for the extent of surgical treatment and postoperative management in children with appendicular peritonitis.

Discussion.

Abdominal operations in children cause postoperative AIO in the unpleasantly high part of cases, with variations depending on the type of initial operation and its clinical peculiarities. According to the modern concept of adhesions pathogenesis development, intraperitoneal inflammation plays an important role in this process [1,2,5]. This correlates with our data [11], indicating that peritonitis is the key condition followed by adhesions formation. It is important to consider as many aspects and variables as feasible to develop clinically relevant and practical predictors. In the present study, we first developed a logistic regression model for adhesive small bowel obstruction prognosis based on the data of a retrospective cohort of 119 pediatric patients.

We also identified, for the first time, the impact of several variables on the possibility of postoperative adhesive intestinal obstruction in patients with appendicular peritonitis using a logistic regression ROC-analysis. These data delineate several key risk factors, associated with adhesions-related sequelae such as type of peritonitis, time spent for the first intervention, incision type, signs of connective tissue dysplasia, leucopenia at the admission, intraabdominal postoperative complications. The absence of vomiting and typical position of the appendix have shown a protective impact.

Understanding which features of operations carry a greater risk of subsequent PAIO will be helpful to prevent acute intestinal obstruction events, as readmission or further surgical interventions. Moreover, our predictive model provides a useful tool for detecting the group of patients that has high chances for adhesion formation and respectively require preventive strategies application on every stage of treatment. The specificity and sensitivity of the developed predictive model account for 77 % and 72 % respectively.

The follow-up evaluation showed that patients who belong to the adhesive obstruction risk group had a considerably higher rate of readmission and requirement for surgical treatment.

Moreover, this cohort of patients has an advanced intraabdominal adhesive process. Our experiences support the active preventive measures in high-risk patients using intraoperative prophylaxis and enhanced recovery after surgery.

Conclusion.

1. Retrospective analysis of 119 case histories determined the level of significance of potential predictors (demographic and individual patient factors, clinical and anamnestic features, laboratory parameters, intraoperative data and volume of surgery, features of the postoperative period course) in children with appendicular peritonitis and PAIO by Pearson's criterion χ^2 . The main factors (leukopenia at hospitalization) ($p = 0.033$) pronounced stigmatization with signs of connective tissue dysplasia ($p = 0.013$) and duration of surgery ($p = 0.0002$) indicated the possibility of the PAIO risk.

2. The method of logistic regression created a prognostic model for calculating the risk of PAIO, which ROC-analysis (test sensitivity – 77%; specificity – 72%) was assessed as “very good”: AUC = 0.796.

3. The proposed prognostic model of the PAIO risk in children with acute inflammatory diseases of the abdominal cavity can be used to optimize treatment strategy at all stages of medical care.

Conflict of interest.

The authors did not find any conflict of interest.

REFERENCES

1. Capella-Monsonís H, Kearns S, Kelly J, et al. Battling adhesions: from understanding to prevention. BMC Biomed Eng. 2019;1:5.
2. Fortin CN, Saed GM, Diamond MP. Predisposing factors to post-operative adhesion development. Hum Reprod Update. 2015;21:536-551.
3. Moris D, Chakedis J, Rahnama-Azar AA, et al. Postoperative Abdominal Adhesions: Clinical Significance and Advances in Prevention and Management. J Gastrointest Surg. 2017;21:1713-1722.
4. Hajibandeh S, Hajibandeh S, Panda N, et al. Operative versus non-operative management of adhesive small bowel obstruction: A systematic review and meta-analysis. Int J Surg. 2017;45:58-66.
5. Melnychenko MH, Kvashnina AA, Vasylychenko LV, et al. Modern viewpoints about pathogenesis of peritoneal adhesions in children. Klinichna khirurgiia. 2016;12:72-75.
6. Deng Y, Wang Y, Guo C. Prediction of surgical management for operated adhesive postoperative small bowel obstruction in a pediatric population. Medicine (Baltimore). 2019;98:e14919.
7. Antomonov MYu. Mathematical processing and analysis of medical and biological data. 2nd edn. Kiev: Medinform. 2017:578.

8. Grigoryev SG, Lobzin YuV, Skripchenko NV. The role and place of logistic regression and roc analysis in solving medical diagnostic task. Zhurnal infektologii. 2016;8:36-45.

9. Koletsi D, Pandis N. Conditional logistic regression. Am J Orthod Dentofacial Orthop. 2017;151:1191-1192.

10. Meurer WJ, Tolles J. Logistic Regression Diagnostics: Understanding How Well a Model Predicts Outcomes. JAMA. 2017;317:1068-1069.

11. Melnichenko MH, Kvashnina AA. Factors of predicting strangulation in children with adhesions to intestinal imperfections. Neonatology, surgery, and perinatal medicine. 2020;4:41-44.

ПРОГНОСТИЧЕСКАЯ МОДЕЛЬ РИСКА ПОСЛЕОПЕРАЦИОННОЙ СПАЕЧНОЙ НЕПРОХОДИМОСТИ КИШЕЧНИКА У ДЕТЕЙ

Мельниченко МГ, Квашнина АА, Ситникова ВА.

Одесский национальный медицинский университет, Одесса, Украина

Резюме.

Цель: разработка прогностической модели риска послеоперационной спаечной непроходимости кишечника (ПСНК) у детей путем использования многомерной статистики.

Материалы и методы. Для создания модели прогнозирования ПСНК путем использования метода бинарной логистической регрессии был проведен ретроспективный анализ особенностей течения острых воспалительных заболеваний органов брюшной полости и лечебной тактики у 119 детей.

Результаты. Был определен уровень значимости клинико-анамнестических признаков у детей с апендикулярным перитонитом и ПСНК, используя критерий Пирсона χ^2 . Выделены основные факторы (лейкопения при госпитализации ($p = 0,033$), выраженная стигматизация признаков соединительнотканной дисплазии ($p = 0,013$) и продолжительность оперативного вмешательства ($p = 0,0002$)), указывающие на возможность риска ПСНК. Методом логистической регрессии создана прогностическая модель для расчета риска ПСНК, которая ROC-анализом (чувствительность теста – 77%; специфичность – 72%) оценивается как «очень хорошая»: AUC = 0,796.

Выводы. Применение предложенной прогностической модели риска ПСНК у детей с острыми воспалительными заболеваниями органов брюшной полости позволяет скорее принять решение относительно объема хирургического лечения и оптимизации лечебной тактики на всех этапах оказания помощи.

Ключевые слова: прогностическая модель, риск, послеоперационная спаечная непроходимость кишечника, дети.